

AMENDMENT TO THE CLAIMS

1. *(Currently Amended)* An additive composition, comprising:
a Mannich reaction product of
 - a) a polyisobutylene alkylated hydroxyaromatic compound;
 - b) an aldehyde; and
 - c) an amine containing at least one reactive amino group,wherein the said polyisobutylene alkylated hydroxyaromatic compound is derived from a combination of a conventional polyisobutylene and a high vinylidene polyisobutylene; and wherein the said polyisobutylene alkylated hydroxyaromatic compound is derived by:
 - i) combining the conventional polyisobutylene and the high vinylidene polyisobutylene prior to the alkylation of the hydroxyaromatic compound; or
 - ii) combining a hydroxyaromatic compound alkylated with the conventional polyisobutylene and a hydroxyaromatic compound alkylated with the high vinylidene polyisobutylene.
2. *(Original)* The additive composition of claim 1 wherein the conventional polyisobutylene has a trisubstituted double bond isomer content of 45 mole % or greater.
3. *(Original)* The additive composition of claim 1 wherein the high vinylidene polyisobutylene has a combined alpha- and beta-vinylidene double bond isomer content of 70 mole % or greater.
4. *(Original)* The additive composition of claim 1 wherein the polyisobutylene of the alkylated hydroxyaromatic compound has an alpha- and beta-vinylidene double bond isomer content of 50 to 95 mole % and a trisubstituted double bond isomer content of 4 to 40 mole %.

5. **(Original)** The additive composition of claim 1 wherein the said polyisobutylene is derived by combining the conventional polyisobutylene and the high vinylidene polyisobutylene prior to the alkylation of the hydroxyaromatic compound.
6. **(Original)** The additive composition of claim 1 wherein the said polyisobutylene is derived by combining a hydroxyaromatic compound alkylated with the conventional polyisobutylene and a hydroxyaromatic compound alkylated with the high vinylidene polyisobutylene.
7. **(Original)** The additive composition of claim 1 wherein the said polyisobutylene is derived by combining a Mannich reaction product from a hydroxyaromatic compound alkylated with the conventional polyisobutylene and a Mannich reaction product from a hydroxyaromatic compound alkylated with the high vinylidene polyisobutylene.
8. **(Original)** The additive composition of claim 1 wherein the said polyisobutylene has a number average molecular weight ranging from 500 to 3,000.
9. **(Original)** The additive composition of claim 1 wherein the hydroxyaromatic compound is phenol, the aldehyde is formaldehyde or a reactive equivalent thereof, and the amine is a secondary monoamine, an alkylenediamine, or a mixture thereof.
10. **(Original)** A fuel additive concentrate composition for an internal combustion engine, comprising:
 - a solvent;
 - the additive composition of claim 1; and
 - optionally one or more additional fuel additives.
11. **(Original)** A fuel composition for an internal combustion engine, comprising:
 - a major amount of a fuel; and
 - a minor amount of the additive composition of claim 1.
12. **(Original)** A fuel composition for an internal combustion engine, comprising:

a major amount of a fuel; and

a minor amount of the fuel additive concentrate composition of claim 10.

13. *(Original)* A method to reduce deposit formation in a fuel system of an internal combustion engine, comprising:

operating the engine with the fuel composition of claim 11.

14. – 16. *(Cancelled)*

17. *(New)* The composition of claim 1 wherein the weight ratio of conventional PIB to high vinylidene PIB is from 25:75 to 60:40.

18. *(New)* The fuel composition of claim 11 wherein the weight ratio of conventional PIB to high vinylidene PIB is from 25:75 to 60:40.

19. *(New)* The method of claim 13 wherein the weight ratio of conventional PIB to high vinylidene PIB is from 25:75 to 60:40.